arXiv:1303.6378v1 [cs.IT] 26 Mar 2013

Cyclic Codes from the Two-Prime Whiteman's Generalized Cyclotomic Sequences with Order 4

Yuhua Sun[1], Tongjiang Yan[2], Hui Li[1]
[1] State Key Laboratory of Integrated Service Networks, Xidian University,
Xi'an, 710071, Shaanxi, China
[2] College of Sciences, China University of Petroleum, Qingdao 266555, Shandong, China
Email: sunyuhua-1@163.com yantoji@163.com lihui@mail.xidian.edu.cn

March 27, 2013

Abstract

Cyclic codes are a subclass of linear codes and have important applications in data storage systems, and communication systems because of their efficient encoding and decoding algorithms. Employing the First Class Whiteman's generalized cyclotomic sequences with order 4, we describe several cyclic codes over the finite field GF(q) and give their generator polynomials. Additionally, Lower bounds on the minimum weight of these cyclic codes are presented.

Index Terms: finite feildcyclotomic sequencecyclic code

1 Introduction

Cyclic codes are a small but highly structured subclass of linear codes. Because of their efficient encoding and decoding algorithms, cyclic codes have wide applications in data storage systems, communication systems. In the past decades years, many important results in the field of cyclic codes have been found, for example, some constructions and properties have been given in [1]-[4]. Recently, Ding [5] presents several classes of cyclic using a class of two-prime Whiteman's generalized cyclotomic sequences in [5], and develops lower bounds on the minimum weight of these cyclic codes. Inspired by the construction of [5] and existing conclusions on another class of Whiteman's generalized cyclotomic sequences with order 4, in this paper, we will give several classes of cyclic codes over the finite field GF(q). Additionally, we present the lower bounds on the minimum weight of these cyclic codes.

2 The linear complexity of sequence and another class of two-prime Whiteman's generalized cyclotomic sequence with order 4

Let q be a power of any prime p, n a positive integer satisfying $\gcd(q, n) = 1$. And let $\mathrm{GF}(q)$ be the finite field with q elements. Assume that $s^n = s_0 s_1 \cdots s_{n-1}$ is a sequence of period n over $\mathrm{GF}(q)$. If there exist n elements $c_1,\ c_2, \cdots, c_l \in \mathrm{GF}(q)$ such that $-s_j = c_1 s_{j-1} + c_2 s_{j-2} + \cdots + c_l s_{j-l},\ j > l$, the sequence s^n is called linear feedback sequence, and the minimal positive integer l is defined as the linear complexity of the sequence s^n. Corespondingly, the polynomial $c(x) = 1 + c_1 x + c_2 x^2 + \cdots + c_l x^l$ is called the feedback polynomial of the sequence s^n. For a periodic sequence, such a positive integer always exists. Any one of feedback polynomial of a sequence is also called a characteristic polynomial of the sequence s^n, and the characteristic polynomial with the minimal length is called the minimal polynomial of the sequence s^n. There are many methods to compute the linear complexity and the minimal polynomial of a periodic sequence. One of these methods is stated in the following lemma.

Lemma 1 *Let s^n be a sequence of period n over $\mathrm{GF}(q)$. Define*

$$s^n(x) = s_0 + s_1 x + s_2 x^2 + \cdots + s_{n-1} x^{n-1} \in \mathrm{GF}(q)[x],$$

then the minimal polynomial $m(x)$ of the sequence s^n is determined by

$$m(x) = \frac{x^n - 1}{\gcd(s^n(x), x^n - 1)}.$$

Consequently, the linear complexity $L(s^n)$ of the sequence s^n is

$$L(s^n) = n - \deg(\gcd(s^n(x), x^n - 1)),$$

where $\deg(f(x))$ represents the degree of the polynomial $f(x)$.

Let p_1, p_2 be two distinct odd primes satisfying $\gcd(p_1 - 1, p_2 - 1) = d$. Let $e = \frac{(p_1-1)(p_2-1)}{d}$ and $n = p_1 p_2$. Let g be a common primitive roots of both p_1 and p_2, and x be an integer satisfying

$$x \equiv g \pmod{p_1},\ x \equiv 1 \pmod{p_2}.$$

Whiteman [6] has proved that

$$\mathbb{Z}_n^* = \{g^s x^i : s = 0, 1, \cdots, e-1;\ i = 0, 1, 2, \cdots, d-1\}.$$

Define

$$W_i = \{g^s x^i : s = 0, 1, \cdots, e-1\},\ i = 0, 1, 2, \cdots, d-1.$$

Then

$$\mathbb{Z}_n^* = \cup_{i=0}^{d-1},\ W_i \cap W_j = \emptyset,\ \text{for } i \neq j,$$

where $\emptyset$ is the empty set. We call W_i two-prime Whiteman generalized class with order d, where $i = 0, 1, 2, \cdots, d-1$.

The cyclotomic numbers corresponding to these cyclotomic classes are defined as

$$(i,j)_d = |(W_i + 1) \cap W_j|,$$

where $0 \leq i, j \leq d-1$ and $W_i + 1 = \{\omega + 1 | \omega \in W_i\}$.

Additionally, for any $r \in \mathbb{Z}_n$, we define

$$d(i,j;r) = |(W_i + r) \cap W_j|,$$

where $W_i + r = \{\omega + r | \omega \in W_i\}$.

Let

$$P_1 = \{p_1, 2p_1, 3p_1, \cdots, (p_2 - 1)p_1\},\ P_2 = \{p_2, 2p_2, 3p_2, \cdots, (p_1 - 1)p_2\}.$$

$$D_0 = \{0\} \cup P_2 \cup \bigcup_{j=0}^{\frac{d}{2}-1} W_j,\ D_1 = P_1 \cup \bigcup_{j=\frac{d}{2}}^{d-1} W_j,$$

$$D_0^* = \{0\} \cup P_2 \cup \bigcup_{j=0}^{\frac{d}{2}-1} W_{2j},\ D_1^* = P_1 \cup \bigcup_{j=0}^{\frac{d}{2}-1} W_{2j+1}.$$

Then the sequence $s^n = (s_i)_{i=0}^{n-1}$ defined by

$$s_i = \begin{cases} 0, & \text{if } i \in D_0, \\ 1, & \text{if } i \in D_1 \end{cases}$$

is called the first class two-prime Whiteman's generalized cyclotomic sequence with order d. And the sequence $s^* = (s_i^*)_{i=0}^{n-1}$ defined by

$$s_i^* = \begin{cases} 0, & \text{if } i \in D_0^*, \\ 1, & \text{if } i \in D_1^* \end{cases}$$

is called the second class two-prime Whiteman's generalized cyclotomic sequence with order d.

For $d = 4$, we have the following lemmas.

Lemma 2 *[7] or [6] Let* $p_1 \equiv p_2 \equiv 1 \pmod 4$ *and* $n = p_1 p_2$, *where* $p_1 \neq p_2$. *There are exactly two representations of* n *in the form* $n = a^2 + 4b^2$ *with* $a \equiv 1 \pmod 4$ *and the sign of* b *indeterminate.*

Let

$$n = a^2 + 4b^2, \quad n = a^{'^2} + 4b^{'^2} \quad (a \equiv a' \equiv 1 \pmod 4) \tag{1}$$

denote those two representations of n in Lemma 2. Then we have

Lemma 3 *[6] Let the symbols be defined as before. Suppose that* $\gcd(p_1-1, p_2-1) = 4$. *Then the sixteen cyclotomic constants* $(i,j)_4$ $(i,j = 0,1,2,3)$ *are associated solely with one of the two decompositions in (1), and the following results hold:*

Case 1. If $\frac{(p_1-1)(p_2-1)}{16}$ *is odd, we have*

$$\begin{aligned}
&(0,0)_4 = \tfrac{1}{8}(3a + 2M + 5),\\
&(0,1)_4 = (1,0)_4 = (3,3)_4 = \tfrac{1}{8}(-a + 4b + 2M + 1),\\
&(0,2)_4 = (2,0)_4 = (2,2)_4 = \tfrac{1}{8}(-a + 2M + 1),\\
&(0,3)_4 = (1,1)_4 = (3,0)_4 = \tfrac{1}{8}(-a - 4b + 2M + 1),\\
&(1,2)_4 = (1,3)_4 = (2,1)_4 = (2,3)_4 = (3,1)_4) = (3,2)_4 = \tfrac{1}{8}(a + 2M - 1).
\end{aligned}$$

Case 2. If $\frac{(p_1-1)(p_2-1)}{16}$ *is even, we have*

$$\begin{aligned}
&(0,0)_4 = (2,0)_4 = (2,2)_4 = \tfrac{1}{8}(-a + 2M + 3),\\
&(0,1)_4 = (1,3)_4 = (3,2)_4 = \tfrac{1}{8}(-a - 4b + 2M - 1),\\
&(0,2)_4 = \tfrac{1}{8}(3a + 2M - 1),\\
&(0,3)_4 = (1,2)_4 = (3,1)_4 = \tfrac{1}{8}(-a - 4b + 2M + 1),\\
&(1,0)_4 = (1,1)_4 = (2,1)_4 = (2,3)_4 = (3,0)_4 = (3,3)_4 = \tfrac{1}{8}(a + 2M + 1),
\end{aligned}$$

where the values of a *and* b *satisfying* $n = p_1p_2 = a^2 + 4b^2$ *depend on the selection of the cyclotomic classes, i.e., the selection of the values* g, *and* $M = \frac{(p_1-2)(p_2-2)-1}{4}$.

Lemma 4 *[8] Let the condition are the same as in Lemma 3. And let* $0 \le i, j \le 3$ *and* $i \ne j$. *Then for any* $r \in P_1 \cup P_2$, *we have*

$$d(i,j;r) = |(W_i + r) \cap W_j| = \frac{(p_1-1)(p_2-1)}{16},$$

and

$$d(i,i;r) = |(W_i + r) \cap W_i| = \begin{cases} \frac{(p_1-1)(p_2-1-4)}{16}, & r \in P_1, \ r \notin P_2, \\ \frac{(p_2-1)(p_1-1-4)}{16}, & r \in P_2, \ r \notin P_1. \end{cases}$$

3 The definition of q-ary cyclic code and its construction

Let p, q, and n be defined as in section 2. And let $\mathrm{GF}(q)^n$ denote the n-dimensional linear space over $\mathrm{GF}(q)$, a linear $[n,k]$ code C over $\mathrm{GF}(q)$ is a k-dimensional subspace of $\mathrm{GF}(q)^n$. If the minimal (Hamming) nonzero weight of C is equal to d_0, we call the linear (n,k) code C a linear $[n,k,d_0]$ code over $\mathrm{GF}(q)$. The linear $[n,k]$ code is called cyclic if $(c_0, c_1, \cdots, c_{n-1}) \in C$ implies $(c_{n-1}, c_0, c_1, \cdots, c_{n-2}) \in C$.

For any vector $(c_0, c_1, \cdots, c_{n-1}) \in \mathrm{GF}(q)^n$, we identify it with the polynomial

$$c_0 + c_1x + c_2x^2 + \cdots + c_{n-1}x^{n-1} \in \mathrm{GF}(q)[x]/(x^n - 1).$$

Then any code C of length n over $\mathrm{GF}(q)$ is equivalent to a subset of $\mathrm{GF}(q)[x]/(x^n - 1)$. It is well known that $\mathrm{GF}(q)[x]/(x^n - 1)$ is a principal ideal ring, i.e., any one of ideals of $\mathrm{GF}(q)[x]/(x^n - 1)$ is principal. Let $C = (g(x))$ be a cyclic code and $h(x) = \frac{x^n-1}{g(x)}$. Then we call $g(x)$ the generator polynomial and $h(x)$ the parity-check polynomial of C.

There are many ways to construct cyclic codes, one class of these cyclic codes is generated by the polynomial

$$\frac{x^n - 1}{\gcd(x^n - 1, S(x))}, \tag{2}$$

where $s^n = (s_i)_{i=0}^{n-1}$ is a sequence of period n over $\mathrm{GF}(q)$ and

$$S(x) = \sum_{i=0}^{n-1} s_i x^i \in \mathrm{GF}(q)[x] \tag{3}$$

is called the generator polynomial of the sequence s^n. Then the cyclic code C_s generated by (2) is called a cyclic code defined by the sequence s^n, which is a linear (n,k) code with $k = \deg(\gcd(x^n - 1, S(x))$, where $\deg(f(x))$ represents the degree of the polynomial $f(x)$. Correspondingly, the sequence s^n is called the defining sequence of the code C_s.

In this paper, we mainly study the cyclic code generated by the first class Whiteman's generalized cyclotomic sequence with order 4.

4 The cyclic codes generated by the first class Whiteman's generalized cyclotomic sequence with order 4

Let p, q, n, p_1, p_2, g, x be defined as before. Without special note, p_1, p_2 always satisfy $\gcd(p_1 - 1, p_2 - 1) = 4$, and $s^n = (s_i)_{i=0}^{n-1}$ always represents the first class Whiteman's generalized cyclotomic sequence with order 4 in this section. Note that $\gcd(n,q) = 1$. Assume that the order of q modulo n

is equal to m. Let α be a primitive element of the finite field $\mathrm{GF}(q^m)$. Then $\beta = \alpha^{\frac{q^m-1}{n}}$ is a primitive nth root of unity.

Define

$$S(x) = \sum_{i \in D_1} x^i = \left(\sum_{i \in P_1} x^i + \sum_{i \in W_2} x^i + \sum_{i \in W_3} x^i \right) x^i \in \mathrm{GF}(q)[x], \tag{4}$$

$$T(x) = \left(\sum_{i \in P_1} x^i + \sum_{i \in W_1} x^i + \sum_{i \in W_2} x^i \right) x^i \in \mathrm{GF}(q)[x], \tag{5}$$

$$R(x) = \sum_{i \in D_1^*} x^i = \left(\sum_{i \in P_1} x^i + \sum_{i \in W_1} x^i + \sum_{i \in W_3} x^i \right) x^i \in \mathrm{GF}(q)[x].$$

Then the polynomial $S(x)$ in (4) is exactly the generator polynomial of the sequence s^n. Our main aim in this section is to find the generator polynomial

$$g(x) = \frac{x^n - 1}{\gcd(x^n - 1, S(x))} \tag{6}$$

of the cyclic code C_s defined by the sequence s^n, where $S(x)$ is the same as in (4). Hence, we need only to find such t's that $S(\beta^t) = 0$ since β is a primitive nth root of unity, where $0 \leq t \leq n-1$. To this end, we need the following lemmas.

Lemma 5 *Let the symbols be defined as above. For any* $r \in W_i$, *we have* $rW_j = W_{i+j \pmod 4}$, *where* $rW_j = \{r\omega | \omega \in W_j\}$.

From the definitions of β and $S(x)$, it is easy to obtain the following result.

Lemma 6 *Let the symbols be defined as above. Then we have*

(I) $\sum\limits_{i \in P_1} \beta^i = \sum\limits_{i \in P_2} \beta^i = -1$;

(II) $\sum\limits_{i \in Z_n^*} \beta^i = \sum\limits_{i \in \bigcup\limits_{j=0}^{3} W_j} \beta^i = 1$;

(III) $S(\beta^0) = S(1) = \frac{(p_1+1)(p_2-1)}{2} \pmod p$.

Proof It is easy to obtain the conclusions in (I) and (III). Now, we will explain the result in (II). Since $\beta^n - 1 = (\beta - 1)(\sum\limits_{i=0}^{n-1} \beta^i) = 0$ and $\beta - 1 \neq 0$, we know that

$$\sum_{i=0}^{n-1} \beta^i = 1 + \sum_{i \in P_1} \beta^i + \sum_{i \in P_2} \beta^i + \sum_{i \in Z_n^*} \beta^i = 0.$$

By (I), we get

$$\sum_{i\in Z_n^*}\beta^i=\sum_{i\in \bigcup\limits_{j=0}^{3} W_j}\beta^i=1+1-1=1.$$

Lemma 7 *Let the symbols be defined as above. For $0\le j\le 3$, we have*

$$\sum_{i\in W_j}\beta^{it}=\begin{cases} -\frac{p_1-1}{4} \pmod p, & \text{if } t\in P_1,\\ -\frac{p_2-1}{4} \pmod p, & \text{if } t\in P_2.\end{cases}$$

Proof Note that for $0\le j\le 3$ we have

$$\begin{aligned} W_j \bmod p_1 &= \{g^s x^j \bmod p_1 :\ s=0,1,2,\cdots,\frac{(p_1-1)(p_2-1)}{4}-1\}\\ &=\{g^{s+j} \bmod p_1 :\ s=0,1,2,\cdots,\frac{(p_1-1)(p_2-1)}{4}-1\}\\ &=\frac{p_2-1}{4}*\{1,2,\cdots,p_1-1\},\end{aligned}$$

where $\frac{p_2-1}{4}$ is the multiplicity of each element in the set $\{1,2,\cdots,p_1-1\}$. Similarly, we have

$$W_j \bmod p_2=\frac{p_1-1}{4}*\{1,2,\cdots,p_2-1\}.$$

Suppose that $t\in P_1$. From Lemma 6 (I), we get

$$\sum_{i\in W_j}\beta^{it}=(\frac{p_1-1}{4})\sum_{i\in P_1}\beta^i=-\left(\frac{p_1-1}{4} \bmod p\right).$$

For $t\in P_2$, we can get the result by similar argument.

Lemma 8 *For all $t\in Z_n$, we have*

$$S(\beta^t)=\begin{cases} -\frac{p_1+1}{2} \pmod p, & \text{if } t\in P_1,\\ \frac{p_2-1}{2} \pmod p, & \text{if } t\in P_2,\\ S(\beta), & \text{if } t\in W_0,\\ -(T(\beta)+1), & \text{if } t\in W_1,\\ -(S(\beta)+1), & \text{if } t\in W_2,\\ T(\beta), & \text{if } t\in W_3,\end{cases}$$

and

$$T(\beta^t) = \begin{cases} -\frac{p_1+1}{2} \pmod p, & \text{if } t \in P_1, \\ \frac{p_2-1}{2} \pmod p, & \text{if } t \in P_2, \\ T(\beta), & \text{if } t \in W_0, \\ S(\beta), & \text{if } t \in W_1, \\ -(T(\beta)+1), & \text{if } t \in W_2, \\ -(S(\beta)+1), & \text{if } t \in W_3. \end{cases}$$

Proof For $t \in P_1$, since $\gcd(p_1, p_2) = 1$, we have

$$tP_1 \bmod n = \{t\omega \bmod n : \ \omega \in P_1\} = P_1.$$

By Lemma 7, we get

$$\begin{aligned} S(\beta^t) = \sum_{i \in D_1} \beta^{ti} &= \left(\sum_{i \in P_1} + \sum_{i \in W_2} + \sum_{i \in W_3} \right) \beta^{ti} \\ &= \sum_{i \in P_1} \beta^i + \sum_{i \in W_2} \beta^{ti} + \sum_{i \in W_3} \beta^{ti} \\ &= -1 - \left(\frac{p_1 - 1}{4} \bmod p \right) - \left(\frac{p_1 - 1}{4} \bmod p \right) \\ &= -\frac{p_1 + 1}{2} \bmod p \end{aligned}$$

For $t \in P_2$, we have

$$tP_1 \bmod n = \{0\}.$$

By Lemma 7 again, we get

$$\begin{aligned} S(\beta^t) = \sum_{i \in D_1} \beta^{ti} &= \left(\sum_{i \in P_1} + \sum_{i \in W_2} + \sum_{i \in W_3} \right) \beta^{ti} \\ &= \sum_{i \in P_1} 1 + \sum_{i \in W_2} \beta^{ti} + \sum_{i \in W_3} \beta^{ti} \\ &= p_2 - 1 - \left(\frac{p_2 - 1}{4} \bmod p \right) - \left(\frac{p_2 - 1}{4} \bmod p \right) \\ &= \frac{p_2 - 1}{2} \bmod p. \end{aligned}$$

If $t \in W_j, 0 \le j \le 3$, by Lemma 5 we have

$$tP_1 \bmod n = P_1, \ tW_i = W_{j+i \bmod 4}.$$

Using Lemma 6 (II) we can get the results.

Corollary 1 *Let the symbols be defined as before. We have the following conclusions.*
(I) If $q \notin W_0$*, we have* $S(\beta) \neq 0, -1$ *and* $T(\beta) \neq 0, -1$.
(II) If $q \in W_0$*, we have* $S^q(\beta) = S(\beta)$ *and* $T^q(\beta) = T(\beta)$.

Proof (I)Note that $\gcd(n, q) = 1$, i.e., $q \in Z_n^*$. If $q \notin W_0$, we know that $q \in \bigcup_{i=1}^{3}$. Without loss of generality, assume that $q \in W_1$. By Lemma 8, we have

$$S^{q^2}(\beta) = S^q(\beta^q) = (-T(\beta) - 1)^q = -T(\beta^q) - 1 = -S(\beta) - 1,$$

i.e.,

$$S^{q^2}(\beta) + S(\beta) + 1 = 0. \tag{7}$$

It is easy to check that none of 0 and -1 is the solution of Eq. (7). Similarly, we have

$$T^{q^2}(\beta) + T^q(\beta) + 1 = 0,$$

i.e., $T(\beta) \neq 0, -1$. For the cases of $q \in W_2$ and $q \in W_2$, the results can be proved by similar argument.

(II) For the case of $q \in W_0$, the conclusion is obvious.

Let $\Omega_i = \frac{p_i + (-1)^{i-1}}{2} \pmod{p}$ and $\Omega = \frac{(p_1+1)(p_2-1)}{2} \bmod p$. Then, from Corollary 1 (I), we have the following theorem.

Theorem 1 *Let the symbols be defined as before. Assume that* $q \notin W_0$*. Then the polynomial* $g(x)$ *in (6) is expressed as*

$$g(x) = \begin{cases} x^n - 1, & \text{if } \Omega_1 \neq 0,\ \Omega_2 \neq 0,\ \Omega \neq 0, \\ \frac{x^n - 1}{x^{p_2} - 1}, & \text{if } \Omega_1 = 0,\ \Omega_2 \neq 0,\ \Omega = 0, \\ \frac{x^n - 1}{x^{p_1} - 1}, & \text{if } \Omega_1 \neq 0,\ \Omega_2 = 0,\ \Omega = 0, \\ \frac{(x^n - 1)(x - 1)}{(x^{p_1} - 1)(x^{p_2} - 1)}, & \text{if } \Omega_1 = \Omega_2 = \Omega = 0. \end{cases}$$

In this case, the cyclic code C_s *over* GF(q) *defined by the first class Whiteman's generalized cyclotomic sequence* s^n *with order 4 has generator polynomial* $g(x)$ *as above.*

Proof Note that $\Omega_i = 0$ results in $\Omega = 0$ and that both $\Omega_1 \neq 0$ and $\Omega_2 \neq 0$ result in $\Omega \neq 0$, where $i = 1, 2$. Then the generator polynomial of the cyclic code C_s defined by s^n follows from (6), Lemma 8, and Corollary 1. □

From (II) in Corollary 1 we know that $S(\beta),\ T(\beta) \in$ GF(q) if $q \in W_0$. Hence, it is possible for $S(\beta) = 0,\ -1$ and $T(\beta) = 0,\ -1$. Next, we will derive the expression of $g(x)$ in (6) for the case of $q \in W_0$. To this end, we need the following lemmas.

Lemma 9 *[6] Let the symbols be defined as before. We denote* $\rho = \frac{(p_1-1)(p_2-1)}{16}$. *Then, for* $-1 \in Z_n^*$, *we have that* $-1 \in W_0$ *if* ρ *is odd and that* $-1 \in W_2$ *if* ρ *is even.*

Lemma 10 *Let the symbols be defined as before. We have the following conclusions.*
(I) $R(\beta)(R(\beta)+1) = \frac{n-1}{4} \pmod{p}$.
(II) If ρ *is odd, we have*

$$S(\beta)(S(\beta)+1) = \frac{n-1}{4} - \frac{b}{2}(2R(\beta)+1),$$

$$T(\beta)(T(\beta)+1) = \frac{n-1}{4} + \frac{b}{2}(2R(\beta)+1).$$

Consequently, we have

$$S(\beta)(S(\beta)+1)T(\beta)(T(\beta)+1) = \left(\left(\frac{n-1}{4}\right)^2 - \frac{b^2}{4}n\right) \pmod{p}.$$

(III) If ρ *is even, we have*

$$S(\beta)(S(\beta)+1) = -\frac{n-1}{4} + \frac{b-1}{2} - b(R(\beta)+1),$$

$$T(\beta)(T(\beta)+1) = -\frac{n-1}{4} - \frac{b+1}{2} + b(R(\beta)+1).$$

Consequently, we have

$$S(\beta)(S(\beta)+1)T(\beta)(T(\beta)+1) = \left(\left(\frac{n-1}{4}\right)^2 - \frac{b^2-1}{4}n\right) \pmod{p},$$

where b *is the same as in Lemma 3.*

Proof (I). In fact, this conclusion is a consequence of Lemma 3.13 in [5]. Here, we omit its proof.
(II). By the definition of $S(x)$ and Lemma 6 (I), we have

$$S(\beta) = -1 + \sum_{i \in W_2} \beta^i + \sum_{i \in W_3} \beta^i.$$

Then we get

$$\begin{aligned} & S(\beta)(S(\beta)+1) \\ = & S(\beta) + \left(-1 + \sum_{i \in W_2} \beta^i + \sum_{i \in W_3} \beta^i\right)^2 \\ = & -\left(\sum_{i \in W_2} \beta^i + \sum_{i \in W_3} \beta^i\right) + \sum_{j \in W_2}\sum_{i \in W_2} \beta^{j+i} + \sum_{j \in W_3}\sum_{i \in W_2} \beta^{j+i} + \sum_{j \in W_2}\sum_{i \in W_3} \beta^{j+i} + \sum_{j \in W_3}\sum_{i \in W_3} \beta^{j+i}. \quad (8) \end{aligned}$$

For odd ρ and $0 \leq j \leq 3$, we know that $-1 \in W_0$ and that $-W_j = \{-\omega : \ \omega \in W_j\} = W_j$ from Lemma 9 and Lemma 5. Furthermore, for any $r \in W_j$, we have $r^{-1} \bmod n \in W_{4-j \bmod 4}$. Hence, we get

$$\sum_{j\in W_2}\sum_{i\in W_2}\beta^{j+i} = \sum_{j\in W_2}\sum_{i\in W_2}\beta^{j-i}$$

$$= |W_2| + \sum_{r\in P_1\cup P_2} d(2,2;r)\beta^r + (2,2)_4\sum_{i\in W_0}\beta^i + (1,1)_4\sum_{i\in W_1}\beta^i + (0,0)_4\sum_{i\in W_2}\beta^i + (3,3)_4\sum_{i\in W_3}\beta^i, \quad (9)$$

$$\sum_{j\in W_2}\sum_{i\in W_3}\beta^{j+i} = \sum_{j\in W_2}\sum_{i\in W_3}\beta^{j-i}$$

$$= \sum_{r\in P_1\cup P_2} d(2,3;r)\beta^r + (2,3)_4\sum_{i\in W_0}\beta^i + (1,2)_4\sum_{i\in W_1}\beta^i + (0,1)_4\sum_{i\in W_2}\beta^i + (3,0)_4\sum_{i\in W_3}\beta^i, \quad (10)$$

$$\sum_{j\in W_3}\sum_{i\in W_2}\beta^{j+i} = \sum_{j\in W_3}\sum_{i\in W_2}\beta^{j-i}$$

$$= \sum_{r\in P_1\cup P_2} d(3,2;r)\beta^r + (3,2)_4\sum_{i\in W_0}\beta^i + (2,1)_4\sum_{i\in W_1}\beta^i + (1,0)_4\sum_{i\in W_2}\beta^i + (0,3)_4\sum_{i\in W_3}\beta^i, \quad (11)$$

$$\sum_{j\in W_3}\sum_{i\in W_3}\beta^{j+i} = \sum_{j\in W_3}\sum_{i\in W_3}\beta^{j-i}$$

$$= |W_3| + \sum_{r\in P_1\cup P_2} d(3,3;r)\beta^r + (3,3)_4\sum_{i\in W_0}\beta^i + (2,2)_4\sum_{i\in W_1}\beta^i + (1,1)_4\sum_{i\in W_2}\beta^i + (0,0)_4\sum_{i\in W_3}\beta^i. \quad (12)$$

Substituting (9-12) into (8) and combining Lemma 3, Lemma 4, Lemma 6 (II), the definition of $R(x)$, and the result of (I) in this lemma, we can get

$$S(\beta)(S(\beta)+1) = \frac{n-1}{4} - \frac{b}{2}(2R(\beta)+1).$$

By the same argument as above, we can get

$$T(\beta)(T(\beta)+1) = \frac{n-1}{4} + \frac{b}{2}(2R(\beta)+1).$$

Using the conclusion of (I) in this lemma, we can get

$$S(\beta)(S(\beta)+1)T(\beta)(T(\beta)+1) = \left(\left(\frac{n-1}{4}\right)^2 - \frac{b^2}{4}n\right) \pmod{p}.$$

(III). Similar to the argument of (II) in this lemma, the results follow.

Remark 1 *For convenience, we will use the following notations.*

(I) For odd $\frac{(p_1-1)(p_2-1)}{16}$*, let*

$$\Omega_3 = \left(\left(\frac{n-1}{4} \right)^2 - \frac{b^2}{4} n \right) \bmod p,$$

$$\Omega_{31} = \frac{n-1}{4} - \frac{b}{2}(2R(\beta)+1),$$

$$\Omega_{32} = \frac{n-1}{4} + \frac{b}{2}(2R(\beta)+1).$$

(II) For even $\frac{(p_1-1)(p_2-1)}{16}$*, let*

$$\Omega_3 = \left(\left(\frac{n-1}{4} \right)^2 - \frac{b^2-1}{4} n \right) \bmod p,$$

$$\Omega_{31} = -\frac{n-1}{4} + \frac{b-1}{2} - b(R(\beta)+1),$$

$$\Omega_{32} = -\frac{n-1}{4} - \frac{b+1}{2} + b(R(\beta)+1).$$

(3) Let $\omega_j(x) = \prod_{i \in W_j} (x - \beta^i)$*. Then we know that* $x^n - 1 = \frac{(x^{p_1}-1)(x^{p_2}-1)}{x-1} \prod_{j=0}^{3} \omega_j(x)$*. Hence we have the following theorem.*

Theorem 2 *Let the symbols be defined as before. Let* $p \neq 2$ *and* $q \in W_0$*. Then we have the following results.*

(I) For $\Omega_1 \neq 0$ *and* $\Omega_2 \neq 0$*, let* $g_1(x)$ *denote the generator polynomial of cyclic code generated by the first class Whiteman's generalized cyclotomic sequence with order 4 in (6). Then we have*

$$g_1(x) = \begin{cases} x^n - 1, & \text{if } \Omega_3 \neq 0, \\ \frac{x^n-1}{\omega_0(x)}, & \text{if } \Omega_{31} = \Omega_3 = 0,\ S(\beta) = 0,\ \Omega_{32} \neq 0, \\ \frac{x^n-1}{\omega_2(x)}, & \text{if } \Omega_{31} = \Omega_3 = 0,\ S(\beta) = -1,\ \Omega_{32} \neq 0, \\ \frac{x^n-1}{\omega_3(x)}, & \text{if } \Omega_{32} = \Omega_3 = 0,\ T(\beta) = 0,\ \Omega_{31} \neq 0, \\ \frac{x^n-1}{\omega_1(x)}, & \text{if } \Omega_{32} = \Omega_3 = 0,\ T(\beta) = -1,\ \Omega_{31} \neq 0, \\ \frac{x^n-1}{\omega_0(x)\omega_3(x)}, & \text{if } \Omega_{31} = \Omega_{32} = \Omega_3 = 0,\ S(\beta) = T(\beta) = 0, \\ \frac{x^n-1}{\omega_0(x)\omega_1(x)}, & \text{if } \Omega_{31} = \Omega_{32} = \Omega_3 = 0,\ S(\beta) = 0,\ T(\beta) = -1, \\ \frac{x^n-1}{\omega_2(x)\omega_3(x)}, & \text{if } \Omega_{31} = \Omega_{32} = \Omega_3 = 0,\ S(\beta) = -1,\ T(\beta) = 0, \\ \frac{x^n-1}{\omega_1(x)\omega_2(x)}, & \text{if } \Omega_{31} = \Omega_{32} = \Omega_3 = 0,\ S(\beta) = T(\beta) = -1. \end{cases}$$

(II) For $\Omega_1 \neq 0$ *and* $\Omega_2 = \Omega = 0$*, let* $g_2(x)$ *denote the generator polynomial of cyclic code generated by the first class Whiteman's generalized cyclotomic sequence with order 4 in (6). And let* $g_1(x)$ *be the*

same as in (I). Then we have $g_2(x)=\frac{g_1(x)}{x^{p_1}-1}$.

(III) For $\Omega_2\neq 0$ *and* $\Omega_1=\Omega=0$, *let* $g_3(x)$ *denote the generator polynomial of cyclic code generated by the first class Whiteman's generalized cyclotomic sequence with order 4 in (6). And let* $g_1(x)$ *be the same as in (I). Then we have* $g_3(x)=\frac{g_1(x)}{x^{p_2}-1}$.

(IV) For $\Omega_1=\Omega_2=\Omega=0$, *let* $g_4(x)$ *denote the generator polynomial of cyclic code generated by the first class Whiteman's generalized cyclotomic sequence with order 4 in (6). And let* $g_1(x)$ *be the same as in (I). Then we have* $g_4(x)=\frac{g_1(x)(x-1)}{(x^{p_1}-1)(x^{p_2}-1)}$.

Proof. Note that $\Omega_{3i}=0$ results in $\Omega_3=0$, where $i=1,2$. And note that $\Omega_{31}=0$ results in $S(\beta)=0$ or $S(\beta)=-1$ and $\Omega_{32}=0$ results in $T(\beta)=0$ or $T(\beta)=-1$. By similar argument to that in Theorem 1 the results follow. □

Corollary 2 *Let the symbols be defined as before. For* $\frac{n-1}{4}\equiv 0 \pmod p$, $p\neq 2$, *and* $q\in W_0$, *then the polynomial* $g_1(x)$ *in Theorem 2 become the following two cases.*

(I) If ρ *is odd, then*

$$g_1(x)=\begin{cases} x^n-1, & \text{if } \frac{b}{2}\not\equiv 0 \pmod p,\\ \frac{x^n-1}{\omega_0(x)\omega_3(x)}, & \text{if } \frac{b}{2}\equiv 0 \pmod p \text{ and } S(\beta)=T(\beta)=0,\\ \frac{x^n-1}{\omega_0(x)\omega_1(x)}, & \text{if } \frac{b}{2}\equiv 0 \pmod p \text{ and } S(\beta)=0,\ T(\beta)=-1,\\ \frac{x^n-1}{\omega_2(x)\omega_3(x)}, & \text{if } \frac{b}{2}\equiv 0 \pmod p \text{ and } S(\beta)=-1,\ T(\beta)=0,\\ \frac{x^n-1}{\omega_1(x)\omega_2(x)}, & \text{if } \frac{b}{2}\equiv 0 \pmod p \text{ and } S(\beta)=T(\beta)=-1. \end{cases}$$

(II) If ρ *is even, let* $R(\beta)=-i,\ i\in\{0,1\}$, *then*

$$g_1(x)=\begin{cases} x^n-1, & \text{if } \frac{b+1}{2}\not\equiv 0 \pmod p \text{ and } \frac{b-1}{2}\not\equiv 0 \pmod p,\\ \frac{x^n-1}{\omega_0(x)}, & \text{if } \frac{b+(-1)^i}{2}\equiv 0 \pmod p \text{ and } S(\beta)=0,\\ \frac{x^n-1}{\omega_2(x)}, & \text{if } \frac{b+(-1)^i}{2}\equiv 0 \pmod p \text{ and } S(\beta)=-1,\\ \frac{x^n-1}{\omega_3(x)}, & \text{if } \frac{b-(-1)^i}{2}\equiv 0 \pmod p \text{ and } T(\beta)=0,\\ \frac{x^n-1}{\omega_1(x)}, & \text{if } \frac{b-(-1)^i}{2}\equiv 0 \pmod p \text{ and } T(\beta)=-1, \end{cases}$$

And for $j=2,3,4$, *the polynomials* $g_j(x)$ *are the same as in Theorem 2.*

Proof. By Lemma 10, we know that $\frac{n-1}{4}\equiv 0 \pmod p$ implies $R(\beta)=0$ or $R(\beta)=-1$. Then the results follow. □

Theorem 3 *Let the symbols be defined as before. Let* $p=2$ *and* $q\in W_0$. *Then we have the following results.*

(I) For odd ρ, the generator polynomial $g(x)$ of cyclic code generated by the first class Whiteman's generalized cyclotomic sequence with order 4 in (6) can be expressed as

$$g(x)=\begin{cases}\frac{x^n-1}{x^{p_2}-1}, & \text{if } b\equiv 2 \pmod 4,\\ \frac{x^n-1}{\omega_0(x)\omega_3(x)(x^{p_2}-1)}, & \text{if } b\equiv 0 \pmod 4 \text{ and } S(\beta)=T(\beta)=0,\\ \frac{x^n-1}{\omega_0(x)\omega_1(x)(x^{p_2}-1)}, & \text{if } b\equiv 0 \pmod 4 \text{ and } S(\beta)=0,\ T(\beta)=1,\\ \frac{x^n-1}{\omega_2(x)\omega_3(x)(x^{p_2}-1)}, & \text{if } b\equiv 0 \pmod 4 \text{ and } S(\beta)=1,\ T(\beta)=0,\\ \frac{x^n-1}{\omega_1(x)\omega_2(x)(x^{p_2}-1)}, & \text{if } b\equiv 0 \pmod 4 \text{ and } S(\beta)=T(\beta)=1.\end{cases}$$

(II) For even ρ, let $R(\beta)=i$, $i=0,1$. Then the generator polynomial $g(x)$ of cyclic code generated by the first class Whiteman's generalized cyclotomic sequence with order 4 in (6) can be expressed as

$$g(x)=\begin{cases}\frac{x^n-1}{(x^{p_2}-1)\omega_0(x)}, & \text{if } b\equiv (-1)^{i+1} \pmod 4 \text{ and } S(\beta)=0,\\ \frac{x^n-1}{(x^{p_2}-1)\omega_2(x)}, & \text{if } b\equiv (-1)^{i+1} \pmod 4 \text{ and } S(\beta)=-1,\\ \frac{x^n-1}{(x^{p_2}-1)\omega_3(x)}, & \text{if } b\equiv (-1)^{i} \pmod 4 \text{ and } T(\beta)=0,\\ \frac{x^n-1}{(x^{p_2}-1)\omega_1(x)}, & \text{if } b\equiv (-1)^{i} \pmod 4 \text{ and } T(\beta)=-1.\end{cases}$$

Proof. First note that $\frac{n-1}{4}\equiv 0 \pmod 2$. For odd ρ, we can derive that b is an even integer, i.e., $b\equiv 2 \pmod 4$ or $b\equiv 0 \pmod 4$. And for even ρ, we can derive that b is odd, i.e., $b\equiv 1 \pmod 4$ or $b\equiv 3 \pmod 4$. By the similar argument to that in Theorem 1, the results follows. □

Remark 2 *When $p=2$ and $p_1\equiv p_2\equiv 1 \bmod 4$, we must have $\frac{n-1}{4}\equiv 0 \bmod p$. By Lemma 3.14 of [5], we know that $2=p\in W_0\cup W_2$ and $q\in W_0\cup W_2$. In fact, for this case, we have the following conjecture.* □

Conjecture 1 *Let $p_1,\ p_2,\ n$ satisfying $\gcd(p_1-1,p_2-1)=4$ be defined as before. Suppose that $\frac{(p_1-1)(p_2-1)}{16}$ is odd. Let $p=2$ and q a power of 2. Then the following results hold.*
(I) There always exist two common primitive roots $g,\ g^*$ of both p_1 and p_2 such that $2\in W_0$ and $2\in W_2^*$, where $W_i=\{g^s x^i : s=0,1,2,\cdots,\frac{(p_1-1)(p_2-1)}{4}-1\}$, $W_i^*=\{g^{*s}x^{*i} : s=0,1,2,\cdots,\frac{(p_1-1)(p_2-1)}{4}-1\}$, $x\equiv g \bmod p_1$ and $x\equiv 1 \bmod p_2$, $x^*\equiv g^* \bmod p_1$ and $x^*\equiv 1 \bmod p_2$, and $i=0,1,2,3$.
(II) Let $(i,j)_4$ and $(i,j)_4^*$ denote the cyclotomic numbers corresponding to the cyclotomic classes $W_{i'}$ and $W_{j'}^*$ respectively in (I), where $0\le i,j,i',j'\le 3$. And let a,b and a^*,b^* denote the two parameters of $(i,j)_4$ and $(i,j)_4^*$ respectively in Lemma 3. Then we have $b\equiv 0 \bmod 4$ and $b^*\equiv 2 \bmod 4$. Consequently, let $g(x)$ and $g^*(x)$ denote the generator polynomial of cyclic code constructed in this paper corresponding

to $W_{i'}$ and $W^*_{j'}$ respectively, then we have

$$g(x) = \begin{cases} \frac{x^n-1}{\omega_0(x)\omega_3(x)(x^{p_2}-1)}, \text{ if } S(\beta)=T(\beta)=0, \\ \frac{x^n-1}{\omega_0(x)\omega_1(x)(x^{p_2}-1)}, \text{ if } S(\beta)=0,\ T(\beta)=1, \\ \frac{x^n-1}{\omega_2(x)\omega_3(x)(x^{p_2}-1)}, \text{ if } S(\beta)=1,\ T(\beta)=0, \\ \frac{x^n-1}{\omega_1(x)\omega_2(x)(x^{p_2}-1)}, \text{ if } S(\beta)=T(\beta)=1 \end{cases}$$

and

$$g^*(x) = \frac{x^n-1}{x^{p_2}-1},$$

where $S(x)$, $T(x)$ denote the polynomials in (4) and (5) corresponding to $W_{i'}$ in (I), and $0 \le i' \le 3$.$\square$

5 The minimal distance of the cyclic codes in this paper

In this section, we computation the minimum distance of some of the cyclic codes and give lower bounds for other cyclic codes in section 4. By the same argument as that in [5], we can get the following two results immediately.

Theorem 4 *Let C_i denote the cyclic code over $\mathrm{GF}(q)$ generated by the polynomial $g_i(x) = \frac{x^n-1}{x^{p_i}-1}$. The C_i has parameters $[n, p_i, d_i]$, where $d_i = p_{i-(-1)^i}$ and $i = 1, 2$.*

Theorem 5 *Let C_{p_1,p_2} denote the cyclic code over $\mathrm{GF}(q)$ generated by the polynomial $g_{p_1,p_2}(x) = \frac{(x^n-1)(x-1)}{(x^{p_1}-1)(x^{p_2}-1)}$. Then C_{p_1,p_2} has parameters $[n, p_1+p_2-1, d_{p_1,p_2}]$, where $d_{p_1,p_2} = \min(p_1, p_2)$.*

We also derive the following lower bounds of the minimum distances of other cyclic codes.

Theorem 6 *Suppose that $q \in W_0$. Let C_i and d_i be defined as in Theorem 4. Let $C_{i,j}$ denote the cyclic code over $\mathrm{GF}(q)$ generated by the polynomial $g_{i,j}(x) = \frac{x^n-1}{(x^{p_i}-1)\omega_j(x)}$. Then $C_{i,j}$ has parameters $[n, p_i + \frac{(p_1-1)(p_2-1)}{4}, d_{i,j}]$, where $d_{i,j} \ge \lceil \sqrt{p_{i-(-1)^i}} \rceil$, $i = 1, 2$, and $0 \le j \le 3$.*

Proof. Note that for $0 \le j \le 3$ and for any $r \in W_j$ we have $r^{-1} \pmod{n} \in W_{4-j \bmod 4}$. Let $i = 1$, $j = 0$ and $c(x) \in \mathrm{GF}(q)[x]/(x^n-1)$ be a codeword of Hanmming weight ω in $C_{1,0}$. Take $r \in W_1$, then $r^{-1} \pmod{n} \in W_3$ and $c(x^r)$ is a codeword of Hamming weight ω in $C_{1,3}$, which implies that $d_{1,0} = d_{1,3}$. By taking $r \in W_2$, $r \in W_3$ respectively, we can get $d_{1,0} = d_{1,2}$ and $d_{1,0} = d_{1,1}$ respectively. Further, for any $j \in \{1, 2, 3\}$ and $r \in W_j$, we have that $c(x)c(x^r)$ is a codeword of C_1. Hence, from Theorem 4, we have $d_{1,j}^2 \ge d_1 = p_2$, i.e., $d_{1,j} \ge \lceil \sqrt{p_2} \rceil$, where $j = 0, 1, 2, 3$. By similar argument, we get $d_{2,j} \ge \lceil \sqrt{p_1} \rceil$, where $j = 0, 1, 2, 3$.

Theorem 7 *Suppose that* $q \in W_0$. *Let* C_{p_1,p_2} *and* d_{p_1,p_2} *be defined as in Theorem 5. The cyclic code over* $\mathrm{GF}(q)$ *generated by the polynomial* $g_{p_1,p_2,j}(x) = \frac{(x^n-1)(x-1)}{(x^{p_1}-1)(x^{p_2}-1)\omega_j(x)}$ *has parameters* $[n, p_1+p_2-1+\frac{(p_1-1)(p_2-1)}{4}, d_{p_1,p_2,j}]$, *where* $d_{p_1,p_2,j} \geq \lceil\sqrt{\min(p_1,p_2)}\rceil$ *and* $0 \leq j \leq 3$.

Proof. Let $j = 0$ and $c(x) \in \mathrm{GF}(q)[x]/(x^n-1)$ be a codeword of Hanmming weight ω in $C_{p_1,p_2,0}$. Take $r \in W_1$, then $r^{-1} \pmod n \in W_3$ and $c(x^r)$ is a codeword of Hamming weight ω in $C_{p_1,p_2,3}$, which implies that $d_{p_1,p_2,0} = d_{p_1,p_2,3}$. By taking $r \in W_2$, $r \in W_3$ respectively, we can get $d_{p_1,p_2,0} = d_{p_1,p_2,2}$ and $d_{p_1,p_2,0} = d_{p_1,p_2,1}$ respectively. Further, for any $j \in \{1,2,3\}$ and $r \in W_j$, we have that $c(x)c(x^r)$ is a codeword of C_{p_1,p_2}. Hence, from Theorem 5, we have $d^2_{p_1,p_2,j} \geq d_{p_1,p_2} = \min(p_1,p_2)$, i.e., $d_{p_1,p_2,j} \geq \lceil\sqrt{\min(p_1,p_2)}\rceil$, where $j = 0,1,2,3$.

Theorem 8 *Suppose that* $q \in W_0$. *Let* C_i *and* d_i *be defined as in Theorem 4. The cyclic code over* $\mathrm{GF}(q)$ *generated by the polynomial* $g_{i,j,h}(x) = \frac{x^n-1}{(x^{p_i}-1)\omega_j(x)\omega_h(x)}$ *has parameters* $[n, p_i + \frac{(p_1-1)(p_2-1)}{2}, d_{i,j,h}]$, *where* $d_{i,j,h} \geq \lceil\sqrt{p_{i-(-1)^i}}\rceil$, $(j,h) \in \{(0,1),(0,3),(1,2),(2,3)\}$, *and* $i = 1,2$.

Proof. Let $i = 1$, $j = 0$, $h = 1$ and $c(x) \in \mathrm{GF}(q)[x]/(x^n-1)$ be a codeword of Hanmming weight ω in $C_{1,0,1}$. Take $r \in W_1$, then $r^{-1} \pmod n \in W_3$ and $c(x^r)$ is a codeword of Hamming weight ω in $C_{1,0,3}$, which implies that $d_{1,0,1} = d_{1,0,3}$. By taking $r \in W_2$, $r \in W_3$ respectively, we can get $d_{1,0,1} = d_{1,2,3}$ and $d_{1,0,1} = d_{1,1,2}$ respectively. Further, for any $r \in W_2$, we have that $c(x)c(x^r)$ is a codeword of C_1. Hence, from Theorem 4, we have $d^2_{1,0,1} \geq d_1 = p_2$, i.e., $d_{1,0,1} \geq \lceil\sqrt{p_2}\rceil$. Hence, $d_{1,j,h} \geq \lceil\sqrt{p_2}\rceil$, where $(j,h) \in \{(0,1),(0,3),(1,2),(2,3)\}$. By similar argument, we get $d_{2,j,h} \geq \lceil\sqrt{p_1}\rceil$, where $(j,h) \in \{(0,1),(0,3),(1,2),(2,3)\}$.

Theorem 9 *Suppose that* $q \in W_0$. *Let* C_{p_1,p_2} *and* d_{p_1,p_2} *be defined as in Theorem 5. The cyclic code over* $\mathrm{GF}(q)$ *generated by the polynomial* $g_{p_1,p_2,j,h}(x) = \frac{(x^n-1)(x-1)}{(x^{p_1}-1)(x^{p_2}-1)\omega_j(x)\omega_h(x)}$ *has parameters* $[n, p_1+p_2-1+\frac{(p_1-1)(p_2-1)}{2}, d_{p_1,p_2,j,h}]$, *where*

$$d_{p_1,p_2,j,h} \geq \lceil\sqrt{\min(p_1,p_2)}\rceil, \text{ and } (j,h) \in \{(0,1),(0,3),(1,2),(2,3)\}.$$

Proof. Let $j = 0$, $h = 1$ and $c(x) \in \mathrm{GF}(q)[x]/(x^n-1)$ be a codeword of Hanmming weight ω in $C_{p_1,p_2,0,1}$. Take $r \in W_1$, then $r^{-1} \pmod n \in W_3$ and $c(x^r)$ is a codeword of Hamming weight ω in $C_{p_1,p_2,0,3}$, which implies that $d_{p_1,p_2,0,1} = d_{p_1,p_2,0,3}$. By taking $r \in W_2$, $r \in W_3$ respectively, we can get $d_{p_1,p_2,0,1} = d_{p_1,p_2,2,3}$ and $d_{p_1,p_2,0,1} = d_{p_1,p_2,1,2}$ respectively. Further, for $r \in W_2$, we have that $c(x)c(x^r)$ is a codeword of C_{p_1,p_2}. Hence, from Theorem 5, we have $d^2_{p_1,p_2,0,1} \geq d_{p_1,p_2} = \min(p_1,p_2)$, i.e., $d_{p_1,p_2,0,1} \geq \lceil\sqrt{\min(p_1,p_2)}\rceil$. Hence, we have $d_{p_1,p_2,j,h} \geq \lceil\sqrt{\min(p_1,p_2)}\rceil$, where $(j,h) \in \{(0,1),(0,3),(1,2),(2,3)\}$.

References

[1] E. Betti and M. Sala, ”A new bound for the minimum distance of a cyclic code from its defining set,” *IEEE Trans. Inf. Theory*, vol.52,no. 8, pp. 3700-3706,2006.

[2] M. v. Eupen and J. H. van Lint, ”On the minimum distance of ternary cyclic codes,” *IEEE Transactions on information theory*, vol. 39, No.2, pp. 409-416, 1993.

[3] W. C. Huffman and V. Pless, Fundamentals of Error-Correcting Codes. Cambridge, U.K: Cambridge Univ. Press, 2003.

[4] J. H. van Lint and R. M. Wilson, ”On the minimum distance of cyclic codes,” *IEEE Trans. Inf. Theory,* vol.32, no. 1, pp.23-40,1986.

[5] C. Ding, ”Cyclic Codes From the Two-Prime Sequences,” *IEEE Trans. Inf. Theory,* vol.58, no. 6, pp.3881-3891, 2012.

[6] A. L. Whiteman, ”A family of difference sets,” *Illinois J. Math.*, vol.6, pp.107-121, 1962.

[7] W. J. Leveque, Topics in number theory, vol. 1, Reading, Massachusetts, Addison-Wesley, 1956.

[8] T. W. Cusick, C. Ding, and A. Renvall, Stream Ciphers and Number Theory, North-Holland Mathematical Library, North-Holland, 1998.